

## Cdc25C Recombinant antibody

**Cat:**B35084D

**Company:** HaoKebio

**Uniprot ID:**P30307

**Applications:** IHC:1:200-1:2000

**Organism:**Rabbit

IHC-Polymer:1:800-1:8000

**Species reactivity:**Human

IHC-TSA:1:1000-1:10000

**Molecular Weight Calculation:** 53 kDa

IF:1:50-1:100

**Observed Molecular Weight:** 60 kDa


WB:1:1000-1:5000

**Background:**

This gene encodes a conserved protein that plays a key role in the regulation of cell division. The encoded protein directs dephosphorylation of cyclin B-bound CDC2 and triggers entry into mitosis. It also suppresses p53-induced growth arrest. Multiple alternatively spliced transcript variants of this gene have been described.

4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit

&amp; Mouse Universal Secondary Antibody, RT, 1h.

**Images:**


Sample: HeLa cells, 4% PFA 12-24h

**Source of Reagents:**

发表[中文论文]请标注:Cdc25C(B35084D)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:Cdc25C(B35084D) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.

**Synonyms:**

CDC25; PPP1R60

**Immunogen:**

Recombinant protein

**Isotype:**

IgG

**Subcellular location:**

Nucleus,Cytoplasm

**Purity:**

Affinity purification

**Form:**

Liquid

**Storage Buffer:**

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.

**Storage:**

Store at -20 °C for one year.

**Experimental procedure:**

Antigen retrieval: Citrate buffer (pH 9.0) , Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody,